




Evidence for a Mass Dependent Forward-Backward Asymmetry in Top Quark Pair Production


D. Amidei, University of Michigan
for
The CDF Collaboration



top quark pair production

- hard scatter cm-frame cross-section

$$\frac{d\sigma}{d\cos\theta} \propto \frac{\alpha_s^2}{q^2} \left[1 + \cos^2\theta^* + (1 - \beta^2) \sin^2\theta^* \right] f(\vec{s})$$



- the final state is specified by $\alpha, q^2, \theta^*, \vec{s}$
- α, q^2 well measured in total σ and $M_{t\bar{t}}$ spectrum. SM like.
- here: production angle
 - dependence on q^2, θ^*
 - asymmetry in production angle with respect to beamline

top quark pair production

- hard scatter cm-frame cross-section

$$\frac{d\sigma}{d\cos\theta} \propto \frac{\alpha_s^2}{q^2} \left[1 + \cos^2\theta^* + (1 - \beta^2)\sin^2\theta^* + \frac{q^2}{q^2 - M^2} \cos\theta^* \right]$$


- the final state is specified by

$$\alpha, q^2, \theta^*, \vec{s}$$

- α, q^2 well measured in total σ and $M_{t\bar{t}}$ spectrum. SM like.


- here: production angle

- dependence on q^2, θ^*
- asymmetry in production angle with respect to beamline



top quark pair production

- hard scatter cm-frame cross-section

$$\frac{d\sigma}{d\cos\theta} \propto \frac{\alpha_s^2}{q^2} \left[1 + \cos^2\theta^* + (1 - \beta^2)\sin^2\theta^* + \frac{q^2}{q^2 - M^2} \cos\theta^* \right]$$


- the final state is specified by

$$\alpha, q^2, \theta^*, \vec{s}$$

- α, q^2 well measured in total σ and $M_{t\bar{t}}$ spectrum. SM like.
- here: production angle
 - dependence on $q^2, \theta^* \longleftrightarrow M_{t\bar{t}}, \Delta y$
 - asymmetry in production angle with respect to beamline
- Measurement of inclusive, Δy , and $M_{t\bar{t}}$ dependent asymmetry

prior measurements

- CDF, 1.9 fb^{-1} , inclusive, corrected to “parton-level”

- tt rest frame $A^{t\bar{t}} = 0.24 \pm 0.14$

- NLO QCD $A^{t\bar{t}} = 0.06 \pm 0.01$

PRL 101, 202001 (2008)

- lab (pp) frame $A^{p\bar{p}} = 0.17 \pm 0.08$

- NLO QCD $A^{p\bar{p}} = 0.04 \pm 0.01$

- D0, inclusive, background subtracted “data-level”

- tt rest frame $A^{t\bar{t}} = 0.12 \pm 0.08$ 0.9 fb^{-1}

PRL 100, 142002 (2008)

- $A^{t\bar{t}} = 0.08 \pm 0.04$ 4.3 fb^{-1}

ICHEP 2010

theoretical interest

- exotic gluons
 - massive chiral color
 - RS gluon
 - color sextets, anti-triplets
- IVB'
 - Z'
 - FV $W'Z'$ t-channel
- FV scalars
- effective Lagrangians
- nice theoretical review by Cao et al. PRD 81,014016, arXiv:1003.3461
- model building must contend with
 - total σ in good agreement with SM
 - $d\sigma/dM_{tt}$ in good agreement with SM


this analysis

- 5.3 fb⁻¹
- standard “lepton+jets” selection, reconstruction
- establish rapidity variables, A_{fb} definitions, in tt frame and lab frame
- models
 - LO
 - QCD charge asymmetry
 - color-octet
- rapidity distributions in the data
- correct the rapidity distributions for
 - backgrounds
 - selection efficiency
 - reconstruction smearing
- to find the model independent A_{fb} to compare to theory
- inclusive in tt and lab frame
- rapidity dependence in tt frame
- M_{tt} dependence in lab frame

top pair production and decay

lepton + jets mode

$$q\bar{q} \rightarrow g \rightarrow t\bar{t} \rightarrow (W^+b)(W^-\bar{b}) \rightarrow (l^+\nu b)(q\bar{q}\bar{b}) \rightarrow l^+ + \cancel{E}_T + 4j + \geq 1 \text{ btag}$$



r-z view

event selection


- high p_t lepton (e/ μ)
 - $E_t/p_t > 20 \text{ GeV (}/c)$
 - $|\eta| < 1.0$
- missing $E_t > 20 \text{ GeV}$
- four jets
 - $E_t > 20 \text{ GeV}$
 - $|\eta| < 2.0$
- at least one b-tagged jet
 - $|\eta| < 1.0$
- 1260 events
- 283 ± 50 non- $t\bar{t}$ background
 - established in precision σ measurement
 - mostly W+jets

top reconstruction

$$l^+ + \cancel{E}_T + 4j + \geq 1 \text{ } b\text{tag} \rightarrow (l^+ \nu b)(q \bar{q} \bar{b}) \rightarrow (W^+ b)(W^- \bar{b}) \rightarrow t \bar{t}$$

- Jet-parton assignment, $p_z(\nu)$ via minimum of simple χ^2
 - Constraints: $M_W = 80.4 \text{ GeV}/c^2$, $M_t = 175 \text{ GeV}/c^2$, $b\text{tag} = b$
 - Float jet p_t within errors

$$\chi^2 = \sum_{lep, jets} \frac{(p_t^{i, meas} - p_t^{i, fit})^2}{\sigma_i^2} + \sum_{j=x, y} \frac{(p_j^{UE, meas} - p_j^{UE, fit})^2}{\sigma_j^2} + \frac{(M_{jj} - M_W)^2}{\Gamma_W^2} + \frac{(M_{l\nu} - M_W)^2}{\Gamma_W^2} + \frac{(M_{bjj} - M_{top})^2}{\Gamma_t^2} + \frac{(M_{bl\nu} - M_{top})^2}{\Gamma_t^2}$$




top reconstruction

$$l^+ + \cancel{E}_T + 4j + \geq 1 \text{ btag} \rightarrow (l^+ \nu b)(q\bar{q}\bar{b}) \rightarrow (W^+ b)(W^- \bar{b}) \rightarrow t\bar{t}$$


- Jet-parton assignment, $p_z(\nu)$ via minimum of simple χ^2
 - Constraints: $M_W = 80.4 \text{ GeV}/c^2$, $M_t = 175 \text{ GeV}/c^2$, btag = b
 - Float jet p_t within errors

$$\chi^2 = \sum_{lep, jets} \frac{(p_t^{i, meas} - p_t^{i, fit})^2}{\sigma_i^2} + \sum_{j=x, y} \frac{(p_j^{UE, meas} - p_j^{UE, fit})^2}{\sigma_j^2} + \frac{(M_{jj} - M_W)^2}{\Gamma_W^2} + \frac{(M_{l\nu} - M_W)^2}{\Gamma_W^2} + \frac{(M_{bjj} - M_{top})^2}{\Gamma_t^2} + \frac{(M_{bl\nu} - M_{top})^2}{\Gamma_t^2}$$

χ^2



$M_{t\bar{t}}$



rapidity : lab frame

- each event has a t_{lep} and t_{had} decay

$$+ q_l \Rightarrow t_{\text{leptonic}} + \bar{t}_{\text{hadronic}}$$

$$- q_l \Rightarrow t_{\text{hadronic}} + \bar{t}_{\text{leptonic}}$$

- and a rapidity for each

$$y_{\text{leptonic}} = y_l$$

$$y_{\text{hadronic}} = y_h$$


- simple rapidity variable in lab frame: y_h

- better measured than y_l
- acceptance out to $|\eta| < 2.0$

- charge tag:


- assign charge with lepton from t_{lep}
- interchange of lepton charge \longleftrightarrow interchange of t and \bar{t}
- If assume CP can combine

$$- q \cdot y_h = y_t^{p\bar{p}}$$



rapidity : tt frame

- a longitudinal boost can change the direction of the top quark
 - A is frame dependent!
 - A^{pp} is diluted by the boost



- a frame invariant variable
 - rapidity difference

$$\Delta y_{t\bar{t}} = q \cdot (y_l - y_h)$$

$$= y_t - y_{\bar{t}}$$

- good : decreased dilution from boost (for NLO QCD effect)

$$A_{FB}^{t\bar{t}} \approx 1.5 \times A_{FB}^{p\bar{p}}$$

- bad: decreased precision

$$\delta \Delta y \approx 0.100$$

- great: ease of interpretation:

$$\Delta y_{t\bar{t}} = 2y_t^{t\bar{t}}$$

→ asymmetry in $\Delta y_{t\bar{t}}$ is equal to asymmetry in top quark production angle in tt rest frame

asymmetries

- lab frame asymmetry in $-qy_h$

$$A_{FB}^{p\bar{p}} = \frac{N(-qy_h > 0) - N(-qy_h < 0)}{N(-qy_h > 0) + N(-qy_h < 0)}$$

$$= \frac{N(y_t^{p\bar{p}} > 0) - N(y_t^{p\bar{p}} < 0)}{N(y_t^{p\bar{p}} > 0) + N(y_t^{p\bar{p}} < 0)}$$

- $t\bar{t}$ rest frame asymmetry in Δy :


$$A_{FB}^{t\bar{t}} = \frac{N(\Delta y > 0) - N(\Delta y < 0)}{N(\Delta y > 0) + N(\Delta y < 0)}$$

$$= \frac{N(y_t^{t\bar{t}} > 0) - N(y_t^{t\bar{t}} < 0)}{N(y_t^{t\bar{t}} > 0) + N(y_t^{t\bar{t}} < 0)}$$

- also of interest: uncharged asymmetries in y_h and $y_l - y_h$


$t\bar{t}$ charge asymmetry in QCD

- Halzen, Hoyer, Kim; Brown, Sadhev, Mikaelian; Kuhn, Rodrigo; Ellis, Dawson, Nason; Almeida, Sterman, Vogelsang; Bowen, Ellis, Rainwater



$t\bar{t}$ frame asymmetries

$$A_{\text{FB}} \sim +10\text{-}12 \% \text{ NLO}$$



$$A_{\text{FB}} \sim -7 \% \text{ NLO}$$

$$A_{\text{FB}} \sim 6 \pm 1.0 \% \quad \text{net}$$

- tests of C in strong interaction confounded by difficulty of jet charge
- reconstructed top pair system has full information on charge flow
 - test C in strong interactions at large q^2


expected QCD asymmetries

- MCFM NLO calculation at “parton level”
- MC@NLO + CDFSIM

model	level	$A^{p\bar{p}}$	$A^{t\bar{t}}$	
MCFM	parton	0.038 ± 0.006	0.058 ± 0.009	
MC@NLO	parton	0.032 ± 0.005	0.052 ± 0.008	truth
MC@NLO	$t\bar{t}$	0.018 ± 0.005	0.024 ± 0.005	sim + reco
MC@NLO	$t\bar{t} + \text{bkg}$	0.001 ± 0.003	0.017 ± 0.004	sim + reco + bkg


- MC@NLO:
 - prediction for data level asymmetry in rest frame is zero!
 - prediction for data level asymmetry in $t\bar{t}$ frame < stat precision (0.028)
- Pythia remains good approximation of SM

inclusive distributions (both lepton charges)



- symmetric!

inclusive distributions (both lepton charges)





- symmetric!

separate by lepton charge

negative leptons

positive leptons



Δy



$$A_{\text{FB}} = -0.048 \pm 0.039$$

$$A_{\text{FB}} = 0.067 \pm 0.040$$

y_h



$$A_{\text{FB}} = 0.076 \pm 0.039$$



$$A_{\text{FB}} = -0.070 \pm 0.040$$

separate by lepton charge

negative leptons

positive leptons

Δy





$A_{\text{FB}} = -0.048 \pm 0.039$

$A_{\text{FB}} = 0.067 \pm 0.040$

It's 1) a charge asymmetry 2) CP conserving

y_h




$A_{\text{FB}} = 0.076 \pm 0.039$

$A_{\text{FB}} = -0.070 \pm 0.040$


combine charges

tt frame



- Combined Δy :
 $A_{\text{FB}} = 0.057 \pm 0.028$
- Compare to mc@nlo
 $A_{\text{FB}} = 0.024$

lab frame



- Combined $-q^*y_h$:
 $A_{\text{FB}} = 0.073 \pm 0.028$
- Compare to mc@nlo
 $A_{\text{FB}} = 0.001$


correct to the parton level

- dN/dy parton level histogram
 - parton level bins j w/ contents P_j
- the top data signal
 - $T_i = S_{ij} \times A_j \times P_j$
- where
 - the A_j are the acceptances for each bin
 - the S_{ij} are the bin-to-bin migration ratios
 - both are estimated with Pythia
- dN/dy data level histogram
 - data level bins i w/ contents D_i
 - Sum of top and bkgrd: $D_i = T_i + B_i$
- to propagate data to parton level:
 - $P_j = A_j^{-1} \times S_{ji}^{-1} \times (D_i - B_i)$
- result is optimized when number of bins = 4


backgrounds

- detailed model for all background components
- fully simulated model samples are reconstructed like data
- asymmetries small (but not zero)

tt rest frame




lab frame




backgrounds

- can be checked in events without b-tags. S:B = 0.3
- data and predictions in good agreement

tt rest frame




lab frame




4-bin measurements

tt rest frame




lab frame




sample	level	$A^{t\bar{t}}$	$A^{p\bar{p}}$
data	data	0.057 ± 0.028	0.073 ± 0.028
MC@NLO	$t\bar{t} + \text{bkg}$	0.017 ± 0.004	0.001 ± 0.003
data	signal	0.075 ± 0.037	0.110 ± 0.039
MC@NLO	$t\bar{t}$	0.024 ± 0.005	0.018 ± 0.005
data	parton	0.158 ± 0.074	0.150 ± 0.055
MC@NLO	parton	0.058 ± 0.009	0.038 ± 0.006

4-bin measurements

tt rest frame



lab frame



sample	level	$A^{t\bar{t}}$
data	data	0.057 ± 0.028
MC@NLO	$t\bar{t} + \text{bkg}$	0.017 ± 0.004
data	signal	0.075 ± 0.037
MC@NLO	$t\bar{t}$	0.024 ± 0.005
data	parton	0.158 ± 0.074
MC@NLO	parton	0.058 ± 0.009

n.b.


D0 signal level 4.3 fb^{-1}

0.08 ± 0.04

we agree!



Asymmetry is a function of Δy and $M_{t\bar{t}}$

- in $t\bar{t}$ frame, QCD asymmetry has linear dependence on Δy , $M_{t\bar{t}}$
- NLO prediction from MCFM



- Δy , $M_{t\bar{t}}$ dependence generally of interest in other theories

$A(\Delta y)$, parton level, data



sample level		$ \Delta y < 1.0$	$ \Delta y \geq 1.0$
data	data	0.021 ± 0.031	0.208 ± 0.062
data	parton	$0.026 \pm 0.104 \pm 0.056$	$0.611 \pm 0.210 \pm 0.147$
MCFM	parton	0.039 ± 0.006	0.123 ± 0.018

Δy vs $M_{t\bar{t}}$





FIG. 10: Top: The Δy — M plane. Each dot represents one event, while the shading is the approximate event probability in the standard PYTHIA based prediction.

$A^{t\bar{t}}(M_{t\bar{t}, i})$

50 GeV bins

100 GeV bins



- How to quantify? Two bins: high and low

color octet model

- need to test methodology on large asymmetry
- model: color octets with axial couplings
- this is a test sample. **not a hypothesis**

- after Ferrario and Rodrigo [arXiv:0906.5541](https://arxiv.org/abs/0906.5541)

- thanks to T. Tait for Madgraph


- If $g_A^q = -g_A^t$ get positive asymmetry

- **Octet A**

- $g_V = 0, |g_A| = 3/2$
 - $M_G = 2.0 \text{ TeV}$
 - $\sigma/\sigma_{\text{sm}} = 1.02$
 - $\sim M_{t\bar{t}}$ spectrum compares to Pythia
 - Model: True $A_{t\bar{t}} = 0.16$ Reco $A_{t\bar{t}} = 0.08$
 - Data: Parton $A_{t\bar{t}} = 0.15$, Reco $A_{t\bar{t}} = 0.06$



- **Octet B**

- MG = 1.8 TeV. asymmetries bigger; $\sigma, M_{t\bar{t}}$ discrepancies bigger





color octet model

- Octet A detail



- $A(M)$





- binned like data, compared to MC@NLO
- reasonable model for the data

the two-bin boundary

- simplest $A(M)$: two bins
- high and low mass
- where to put boundary?
- look at significance at high mass vs boundary
- best boundary: 450 GeV/ c^2



bin-edge (GeV/ c^2)	OctetA		OctetB	
	A^{tt}	significance	A^{tt}	significance
345	0.082 ± 0.028	2.90	0.168 ± 0.028	5.99
400	0.128 ± 0.036	3.55	0.235 ± 0.035	6.74
➔ 450	0.183 ± 0.047	3.91	0.310 ± 0.044	7.08
500	0.215 ± 0.060	3.60	0.369 ± 0.054	6.81
550	0.246 ± 0.076	3.25	0.425 ± 0.066	6.43
600	0.290 ± 0.097	2.97	0.460 ± 0.081	5.70

data: Δy at low and high mass





selection	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
reco data	0.057 ± 0.028	-0.016 ± 0.034	0.212 ± 0.049
MC@NLO	0.017 ± 0.004	0.012 ± 0.006	0.030 ± 0.007

Δy at high mass by lepton charge



selection	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
A_{lh}^+	0.067 ± 0.040	-0.013 ± 0.050	0.210 ± 0.066
A_{lh}^-	-0.048 ± 0.039	0.020 ± 0.047	-0.210 ± 0.071

Δy at high mass by lepton charge



selection	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
A_{lh}^+	0.067 ± 0.040	-0.013 ± 0.050	0.210 ± 0.066
A_{lh}^-	-0.048 ± 0.039	0.020 ± 0.047	-0.210 ± 0.071

- consistent with CP conservation
- argues against experimental artifact, as detection/reconstruction are sign independent

unfold to the parton level

- dN/dy parton level histogram
 - parton level bins j w/ contents P_j
- the top data signal
 - $T_i = S_{ij} \times A_j \times P_j$
- where
 - the A_j are the acceptances for each bin
 - the S_{ij} are the bin-to-bin migration ratios
 - both measured with symmetric Pythia
- dN/dy data level histogram
 - parton level bins j w/ contents P_j
 - data: in bins i w/ contents $D_i = T_i + B_i$
- to propagate data to parton level:
 - $P_j = A_j^{-1} \times S_{ji}^{-1} \times (D_i - B_i)$
- result is optimized when number of bins = 4

BUT NOW:

4 bins in Δy and M_{tt}

low mass forward
low mass backward
high mass forward
high mass backward

tests of unfold procedure

Sample	$A^{t\bar{t}}$ level	$M_{t\bar{t}} < 450 \text{ GeV}/c^2$	$M_{t\bar{t}} \geq 450 \text{ GeV}/c^2$
PYTHIA	MC truth	0.002	0.001
	reconstructed	-0.011 ± 0.006	-0.013 ± 0.008
	corrected	0.001 ± 0.018	0.006 ± 0.014
MC@NLO	MC truth	0.043	0.070
	reconstructed	0.015 ± 0.006	0.043 ± 0.009
	corrected	0.066 ± 0.014	0.086 ± 0.011
Octet A	MC truth	0.081	0.276
	reconstructed	0.024 ± 0.035	0.183 ± 0.010
	corrected	0.054 ± 0.022	0.308 ± 0.016
Octet B	MC truth	0.150	0.466
	reconstructed	0.078 ± 0.036	0.310 ± 0.009
	corrected	0.187 ± 0.024	0.476 ± 0.015

sys uncertainty of unfold procedure

Source	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
background size	0.017	0.032
background shape	0.003	0.003
JES	0.005	0.012
ISR/FSR	0.012	0.008
color reconnection	0.009	0.004
PDF	0.018	0.004
physics model	0.035	0.035
total	0.047	0.049


TABLE XII: Systematic uncertainties in the two-mass bin unfold

$A^{t\bar{t}}$ at high and low mass: data, signal, parton level

selection	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
data	-0.016 ± 0.034	0.210 ± 0.049
MC@NLO $t\bar{t}$ +bkg	$+0.012 \pm 0.006$	0.030 ± 0.007
data signal	$-0.022 \pm 0.039 \pm 0.017$	$0.266 \pm 0.053 \pm 0.032$
MC@NLO $t\bar{t}$	$+0.015 \pm 0.006$	0.043 ± 0.009
data parton	$-0.116 \pm 0.146 \pm 0.047$	$0.475 \pm 0.101 \pm 0.049$
MCFM	$+0.040 \pm 0.006$	0.088 ± 0.013

TABLE XIII: Asymmetry $A^{t\bar{t}}$ at high and low mass compared to prediction.

$A^{t\bar{t}}$ at high and low mass: parton level



Studies of A^{tt} at the data level

selection	N events	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
standard	1260	0.057 ± 0.028	-0.016 ± 0.034	0.212 ± 0.049
electrons	735	0.026 ± 0.037	-0.020 ± 0.045	0.120 ± 0.063
muons	525	0.105 ± 0.043	-0.012 ± 0.054	0.348 ± 0.080
data $\chi^2 < 3.0$	338	0.030 ± 0.054	-0.033 ± 0.065	0.180 ± 0.099
data no-b-fit	1260	0.062 ± 0.028	0.006 ± 0.034	0.190 ± 0.050
data single b-tag	979	0.058 ± 0.031	-0.015 ± 0.038	0.224 ± 0.056
data double b-tag	281	0.053 ± 0.059	-0.023 ± 0.076	0.178 ± 0.095
data anti-tag	3019	0.033 ± 0.018	0.029 ± 0.021	0.044 ± 0.035
pred anti-tag	-	0.010 ± 0.007	0.013 ± 0.008	0.001 ± 0.014
pre-tag	4279	0.040 ± 0.015	0.017 ± 0.018	0.100 ± 0.029
pre-tag no-b-fit	4279	0.042 ± 0.015	0.023 ± 0.018	0.092 ± 0.029

Jet multiplicity dependence

- the NLO QCD asymmetry has a strong N_{jet} dependence

selection	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
inclusive	0.024 ± 0.004	0.015 ± 0.005	0.043 ± 0.007
4-jet	0.048 ± 0.005	0.033 ± 0.006	0.078 ± 0.009
5-jet	-0.035 ± 0.007	-0.032 ± 0.009	-0.040 ± 0.012

- data: the high mass asymmetry is significantly reduced for 5 jet events

selection	N events	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
data 4-jet	939	0.065 ± 0.033	-0.023 ± 0.039	0.26 ± 0.057
data 5-jet	321	0.034 ± 0.056	0.0049 ± 0.07	0.086 ± 0.093

- need to study other models, color flow, asymmetry reco in ttj

Frame dependence

- a selection of cross-checks in the lab frame using $-qy_h = y_t^{p\bar{p}}$

selection	all M	$M < 450 \text{ GeV}/c^2$	$M \geq 450 \text{ GeV}/c^2$
data reco	0.073 ± 0.028	0.059 ± 0.034	0.103 ± 0.049
MC@NLO	0.017 ± 0.004	-0.008 ± 0.005	0.022 ± 0.007
A_h^+	-0.076 ± 0.039	-0.085 ± 0.047	-0.053 ± 0.072
A_h^-	0.070 ± 0.040	0.028 ± 0.050	0.148 ± 0.066
single b-tags	0.095 ± 0.032	0.079 ± 0.034	0.130 ± 0.057
double b-tags	-0.004 ± 0.060	-0.023 ± 0.076	0.028 ± 0.097

- the high mass asymmetry is less significant in the lab frame
 - like QCD ?
- the high mass double tag asymmetry is low in the lab frame
 - statistics?
 - $|\eta| < 1.0$ for b-tags. acceptance + physics?

Summary

- Inclusive A in lab and tt frames in 2 sigma excess over SM
- Consistent with CP conservation

- A^{tt} has a strong dependence on Δy , M_{tt}

- For $M_{tt} > 450 \text{ GeV}/c^2$

$$A_{\text{reco}}^{tt} = 0.210 \pm 0.049, \quad A_{\text{parton}}^{tt} = 0.475 \pm 0.112$$
$$A_{\text{NLO reco}}^{tt} = 0.043 \pm 0.006 \quad A_{\text{MCFM}}^{tt} = 0.088 \pm 0.013$$

- The asymmetry at high mass is consistent with CP conservation
- Most cross-checks rule out non-physics, although a few puzzles

- The modest inclusive asymmetry originates with a significant effect at large Δy , M_{tt}

- There is a lot more work to do!